

**Table 1. Number, median days, incidence rate<sup>1</sup> and relative standard errors of nonfatal occupational injuries and illnesses with days away from work<sup>2</sup> involving musculoskeletal disorders<sup>3</sup> by selected natures of injury or illness, Maryland, 2002**

Nature	Number	Median days away from work	Incidence rate	Relative standard error
Total	7,625	6	45.6	5.0
021 Sprains, strains, tears	6,913	6	41.3	5.1
0972 Back pain, hurt back	99	4	0.6	23.6
0973 Soreness, pain, hurt, except the back	97	3	0.6	23.9
1241 Carpal tunnel syndrome	108	53	0.7	22.7
153 Hernia	336	30	2.0	13.3
1531 Inguinal hernia	305	30	1.8	13.9
1539 Hernia, n.e.c.	--	--	--	--
17 Musculoskeletal system and connective tissue diseases and disorders	72	31	0.4	27.7
173 Rheumatism, except the back	72	31	0.4	27.7
1731 Bursitis	--	--	--	--
1733 Tendonitis	47	31	0.3	33.9
1735 Ganglion/cystic tumor	--	--	--	--
1739 Rheumatism, except the back, n.e.c.	18	108	0.1	54.8

<sup>1</sup> Incidence rates represent the number of injuries and illnesses per 10,000 full-time workers and were calculated as: (N / EH) X 20,000,000 where,

N = number of injuries and illnesses,  
EH = total hours worked by all employees during the calendar year,  
20,000,000 = base for 10,000 full-time equivalent workers (working 40 hours per week, 50 weeks per year).

<sup>2</sup> Days away from work include those which result in days away from work with or without job transfer or restriction.

<sup>3</sup> Includes cases where the nature of injury is: sprains, strains, tears; back pain, hurt back; soreness, pain, hurt, except back; carpal tunnel syndrome; hernia; or musculoskeletal system and connective tissue diseases and disorders and when the event or exposure leading to the injury or illness is: bodily reaction/bending, climbing, crawling, reaching, twisting; overexertion; or repetition. Cases of Raynaud's phenomenon, tarsal tunnel syndrome, and herniated spinal discs are not included. Although these cases may be considered MSD's, the survey classifies these cases in categories that also include non-MSD cases.

NOTE: Dashes indicate data that do not meet publication guidelines or data for incidence rates less than .05 per 10,000 full-time workers. The scientifically selected probability sample used was one of many possible samples, each of which could have produced different estimates. A measure of sampling variability for each estimate is available upon request.

SOURCE: Bureau of Labor Statistics, U.S. Department of Labor, April 2004